

10/pets

10/540615

JC20 Rec'd PCT/PTO 23 JUN 2005

RECOMBINANT HEPATITIS A VIRUS ANTIGENS OBTAINED IN PLANT CELLS.

Field of the invention.

This invention is related with the biotechnology branch and more specifically with the expression of recombinant proteins in transgenic plant cells and the use of these plants as antigen vaccines. In particular, it is shown recombinant antigens of hepatitis A virus obtained in transgenic plants derived from the expression of modified fragments of the genome of HAV from the M2 strain isolated in Cuba.

It is also demonstrated the usefulness of these antigens to develop immune response in animals after inoculation by different ways.

Previous art.

The genome of the HAV is a simple strain RNA with positive polarity. It has approximately 7.5 kb codified for a 253 kDa polyprotein (Cohen *et al.*, Journal of Virology (1987), 61:3035-3039). The polyprotein suffers both translational and post-translational processes, creating structural mature proteins (VP1, VP2, VP3, VP4 and 2A) and no structural proteins (2B, 2C, 3A, 3B, 3C and 3D).

The protease 3C ($P_{ro}3C$), present in the P3 domain of the viral polypotein, is the protease which takes part in the cleavage of the polyprotein of HAV (Martín *et al.*, J Virol. (1999), 73(8):6220-7), allowing the liberation of the intermediates P1-2A, 2BC and P3, which are processed afterwards. Therefore, for adequate formation of the envelop and for the replication of the HAV it is necessary the occurrence of a differential proteolytic process of the HAV polyprotein. During the processing of the P3 region, just the site 3C/3D is efficiently cleaved. There is a retarded processing in the sites 3A/3B and/or 3B/3C, allowing the accumulation of the intermediary polypeptide 3ABC (Kusov *et al.*, Journal of Virology (1999), 73:9867-9878), which cleaves the polypeptide P1-2A with similar efficiency of the protease 3C. This step facilitates more efficiency in the pentamers formation. The characteristic form of the virus comes from the unification of the viral proteins and its three dimensional configuration is important for the generation of a protective immune response. The virion of the HAV shows an immunodominant antigenic site of neutralization, which is strictly preserved among strains of HAV isolated from different geographic areas. It is arranged by five conformational epitopes, three of them in the pentamers and the two others which are created after the assembling of the pentamers to form the envelope..

It is considered that these latest epitopes are formed due to conformational changes at the antigenic site or due to juxtaposition of epitope fragments present in the pentamers during

their assembly. Both pentamers as well as the viral particles induce neutralization antibodies and therefore they can be useful for vaccine development (Stapleton *et al.*, Journal of Virology (1993), 67:1080-1085). Using recombinant Baculoviruses containing the complete Open Reading Frame of the HAV, a giant polyprotein of the HAV was expressed. Other intermediary proteins as result of its processing in insect cells were also expressed (Stapleton
5 *et al.*, The Journal of Infectious Diseases (1995), 171:9-14). In addition, recombinants vaccinia viruses were constructed which expressed the same polyprotein of the HAV in mammalian cells. Extracts of cells infected with these genetic constructions showed a post-translational processing of the polyprotein rendering capsides similar to those of the HAV
10 (Winokur *et al.*, Journal of Virology (1994), 65:5029-5036). There are patents describing variants of recombinant vaccines against HAV expressed in baculovirus systems and vaccinias, such as: Publication of Patent Application No WO9301279 Winokur *et al.* On January 21, 1993.; patent No US5294548 (McLinden *et al.*, March, 1994); Publication of Patent Application WO09844122 (Probst, August 27, 2002); Publication of Patent
15 Application WO9111460 and patent US5605692 (Thomas *et al.* On February 25, 1997), where the sequences of the Open Reading Frame (ORF) are used for the production of immunogenic capsids and pentamers, and the methods for the obtention of the capsid of the HAV, expressing the structural regions and the region P3 in orientation cis, trans as well as bicistronic constructions are protected.

20 **Transgenic plant as bioreactors.**

The first transgenic plants originated from gene transference by the rhizobacterium *Agrobacterium tumefaciens* were produced at the beginning of 80's decade (Zambryski *et al.*, EMBO J. 1983, 2: 2143-2150). This technology was initially used as a way to achieve resistance to pathogenic microorganisms (Powell *et al.*, Science 1986, 232: 738-743), to
25 insects (Vaeck *et al.*, Nature 1987, 328: 33-37), and to herbicides (Of Block *et al.*, 1987, EMBO J. 6: 2513-2518). But the demonstration of the capacity of the vegetable cells to ensemble correctly foreign proteins of a high structural complexity rapidly indicated its possible value as a new strategy for escalating the economic production of recombinant proteins of industrial and biopharmaceutical interest (Barta *et al.*, Plant Mol. Biol 1986, 6:
30 347-357; Cramer *et al.*, Ann. N And Acad. Sci. 1996, 792: 62-71; Staub *et al.*, Nature Biotechn. 2001. 18: 333-338).

In 1992, a new concept related to the production of subunits vaccines was introduced. It came from the demonstration that the transgenic plants could express the surface antigen of hepatitis B (HBsAg). Based on these findings it was thought that plants might be used to

produce vaccine candidates in edible products and to achieve immunization only by the consumption of these products. By these facts, the denomination of "**edible vaccines**" appeared (Arntzen *et al.*, Plants. Vaccine 1994, 94:339-344). Later it was demonstrated that mice fed with transgenic potatoes containing the HBsAg showed a primary immune response similar to the one obtained when a unique dose of a commercial vaccine is administered by intraperitoneal way. These results indicated that the expression of antigens in edible plants tissues can be considered as a new route of immunization (Richter *et al.*, Nature Biotechnology 2000, 18:1167-1171).

There are several patents describing the use of plants for the expression of vaccines, such as: patent No US5484719 (Lam *et al.*, January 16, 1996; patent No US5612487 (Lam *et al.*, January 16, 1996); patent No US5914123, divisional of the patent No US5612487 and a continuation in part of the Patent Application No PCT/US94/02332 (Arntzen *et al.*, June 22, 1999); patent No US6136320 (Arntzen *et al.*, June 22, 1999); Publication of Patent Application WO9612801 (Arntzen *et al.*, May 28, 2002) and the Patent Application No US2002006411 (Lam *et al.*, On June 4, 2002).

The documents previously mentioned describe the use of the plants as vaccines, as well as the expression of the HBsAg in plants and in some cases they use the term "viral hepatitis" to refer to the virus of the hepatitis B (VHB). The VHB differs significantly from the virus of hepatitis A with very different characteristics and therefore they belong to different genres from the taxonomic point of view. To achieve recombinant proteins of HAV capable of raising an immunological important response it is necessary to express several proteins of the viral genome and then to achieve that such particles are formed as pentamers or empty capsids. The processing and the formation of immunogenic particles have been achieved only in eukariotic systems as vaccinias and baculoviruses, but no in simpler systems such as yeasts. In transgenic plants, complex antigens as HAV have not been expressed. In the case of the VHB, the antigen is formed by only one protein which is efficiently particulated in simple eucariotic systems such as yeast. Due to the previously exposed arguments we believe that the expression of the HBsAg does not include the expression of pentamers or empty capsides of the HAV in plants. In other patent applications is specifically described the expression of different viral antigens, such as antigens of the virus of the human papilloma in the patent applications No WO0161022 of Sohn *et al.*, of August 23, 2001; that of the virus of the aftosa fever in the patent applications No CN1319670 of Zhong *et al.*, of October 31, 2001; of the rotavirus in the patent applications WO0159070 of Reads *et al.*, of August 16,

2001 and that of the gumboro virus in the patent applications No WO0197839 of Shachar *et al.*, of December 27, 2001.

The production of recombinant proteins in plants offers many potential advantages to generate pharmaceutical compounds or vaccines of importance in clinical medicine. First, the plant systems are more economic than the industrial infrastructure used in fermentation systems or in bioreactors. Secondly, the technology to harvest and to try plants and its products to industrial scale is already available. In third position, the requirement of purification of the compound can be eliminated when the plant that contains recombinant protein is used as food (as in case of the edible vaccines). In fourth place, it is possible to direct the recombinant proteins to certain intracellular compartments as mitochondria, vacuoles, chloroplasts and endoplasmatic reticule, or to express them directly in these compartments (for example in chloroplasts). In fifth place, the risks to the health for possible pollution of the recombinant product with human beings pathogens are minimal. Finally, plants as expression systems of the recombinant proteins of pharmaceutical importance have as additional advantages the fact that many of the steps of the secretory route, including folding, assembly, glycosylation at the level of endoplasmatic reticule are similar to the cells of mammals (Ma y Hein, Plant Physiol. 1995, 109: 341- 346; Rayon *et al.*, J. Exp. Bot. 1998, 49: 1463-1472; Sanderfoot y Raikhel, Plant Cell. 1999, 11: 629-641; Vitale and Denecke, Plant Cell 1999, 11: 615-628; Lerouge *et al.*, Pharmaceutical Biotechnology 2000, 1: 347-354).

Detailed description of the invention.

Basic design of the object of this invention is supported by genetic constructions that allow the combined expression of genes that codify for different variants of structural proteins and mutated non-structural regions, directed to the recombinant expression of antigenic pentamers and capsids of HAV in transgenic plants capable of generating an immune response.

The novelty of our invention fundamentally falls in the regions of the viral genome used for the conformation of a new Open Reading Frame coding for a polyprotein of minor size, shaped by strictly structural regions (only up to the protein 2A) and the viral modified protease which presents a superior size to the protease 3C of the virus due to the fact that the sites of cleavage between the proteins 3A/3B and 3B/3C are mutated. The expression in the cytosol and in the endoplasmatic reticulum of viral capsids of the HAV in transgenic plants is achieved by the first time, specially under the control of the promoters and regulatory signals used. The pentamer and capsids formation in endoplasmatic reticulum as a product of the

combined expression of the structural region and the region responsible for the proteolysis, demonstrates the possibilities of this compartment for the assembly and the storage of complex structures, such as the case of the HAV. The production of pentamers and capsids in plants allows us the possibility to employ them as bioreactors to obtain a cheap and sure vaccine.

The invention is demonstrated by means of examples in which transgenic tobacco, rice and carrot plants were used for the obtaining by the first time of immunogenic capsids and pentamers of the virus in vegetable cells. The capsids and pentamers of HAV, resultant of the invention, can be used as antigens vaccines and also in diagnostic assays for HAV detection.

10 **Genetic constructions.**

Obtaining cDNA of the HAV.

From the RNA of the strain M2 of the HAV isolated in Cuba, the nucleotide sequence coding for the Open Reading Frame (ORF) of the virus was amplified, using Reverse Transcription Technology – Polymerase Chain Reaction (RT-PCR). This fragment was cloned in a plasmid and its nucleotide sequence was determined. It presents differences that generate variations in 11 aminoacid residues with respect to the reported sequences. The analysis of the sequences allows to classify the strain M2 as part of the subgenotype IA, to which belong almost all of the American strains. From the genome of this strain, modified fragments were designed and constructed and then were used in different genetic constructions that are object of this invention,

Genetic constructions of vectors for the expression of capsids and pentamers in transgenic plants.

Recombinant protease of HAV.

To allow the formation of viral capsids it is necessary that a differentially proteolytic processing of the polyprotein happens which allows the ordered releasing of viral proteins. The efficiency of capsid formation increases when the intermediary 3ABC is present due to the hydrophobic interaction of the protein 3AB with the membrane and the viral proteins. The cleavage sites of the protease 3C between the proteins 3A/3B and 3A/3C were mutated to obtain a polypeptide 3ABC from which the protease 3C is not released and in turn preserves its proteolytic function, necessary for formation of the pentamers and capsids of the HAV,. Substitutions were glutamic by valine between 3A/3B and serine by leucine between 3B/3C. This polypeptide was used in the design of novel and different strategies for the expression of HAV proteins forming immunogenic capsids and pentamers.

Recombinant HAV for the expression of capsids and pentamers in the cytosol of the vegetable cell.

In HAV, the polypeptide P1-2A has an important function in the formation of the viral capsid. In this polypeptide there are two signs that regulate the formation of the capsid. In its
 5 carboxil terminal domain is found the protein 2A, which is needed in the first stages of capsid assembling to achieve the formation of pentamers. They are formed from the combination of five not processed molecules of the polypeptide P1-2A. The protein VP4 is needed in the second stage for the association of the pentamers and the formation of the capsids.

For the expression of the modified polyprotein in the citosol of the vegetable cell, there were
 10 constructed vectors that contain the sequence of modified Open Reading Frame (ORFm). These constructs codifies for a significantly minor size polyprotein (compared to the original polyprotein of the HAV). This sequence is the result of the fusion of P1-2A polypeptide coding sequence and the sequence that codifies for the mutated protease 3ABC.

The plasmid vector used for the transformation of plants, by means of *A. tumefaciens*,
 15 contains DNA sequences that codifies for proteins of the HAV fused to the nucleotides sequences coding for regulatory signals of their expression in plants. In this case, the sequence coding for the protein is not fused to any specific signal of transport across the secretory route of the vegetable cell, so it is expressed in the cytosol of the cell.

Recombinant HAV for exclusive expression of pentamers in the cytosol of the vegetable cell.

Autoprocess and form exclusively immunogenic viral pentamers. The minor size of the pentamers regarding the capsids allows to achieve higher levels of expression because it provokes a minor As it has been previously described, the protein VP4 as part
 25 of the polypeptide VP0 is required for the association of the pentamers and the formation of the viral capsids.

From the nucleotide sequence coding for the polyprotein ORFm, the fragment that codes for the protein VP4 was eliminated, giving place to a sequence called Δ ORFm. A plasmid vector for the expression of the polyprotein Δ ORFm fused to the sequences that regulate its expression in the cytosol of the vegetable cells was constructed. It was used to obtain
 30 transgenic plants by means of the infection of leaves of tobacco, rice and carrot with *A. tumefaciens*. The polyprotein expressed from this genetic construction has a significantly minor size and it is capable to metabolic charge in the vegetable cell. The obtained product is equally feasible to be used like immunogen for the development of vaccines.

Recombinant HAV for the expression of capsids and pentamers in the endoplasmic reticulum of the vegetable cell.

The accumulation of heterologous proteins in the endoplasmic reticulum in plants is achieved by means of the use of sequences to drive them to the secretory pathway, and therefore the endoplasmic reticulum, and also of signals of retention in this organelle.

As signal peptide, a sequence that codes for the N-terminal peptide of the sweet potato sporamin was used. As signal of retention of proteins in the endoplasmic reticulum, the sequence that codes for the peptide KDEL located in the carboxyl terminal of the protein, was used.

- 10 Sweet potato sporamin signal peptide was fused to the 5' region of P1-2A nucleotide coding sequence and a sequence for spacer peptide fused to KDEL coding sequence was inserted in the 3' region. Resulting DNA fragment was placed under regulation of plant expression signals at a binary vector. This vector included the mutated polypeptide 3ABC coding sequence fused at its 5' extreme to sweet potato signal peptide and at its 3' extreme the KDEL coding sequence. All this elements also under regulation of plant expression signals. Both polypeptides are located in the endoplasmic reticulum and the protease 3ABC is able to process the polypeptide P1-2A and achieve efficient particle formation.

Recombinant HAV to exclusively express pentamers in the endoplasmic reticulum of the vegetable cell.

- 20 The polypeptide Δ P1-2A was obtained by removing the VP4 nucleotide coding sequence from the P1-2A polypeptide coding sequence. This sequence was fused at its 5' end to sweet potato sporamin signal peptide and at its 3' end, to the sequence that codifies for a spacer peptide joined to KDEL coding sequence. In the same binary vector the sequence that codes for the mutated polypeptide 3ABC was equally fused at its 5' end to the sweet potato sporamin signal peptide and at its 3' end to the KDEL coding sequence. All the elements under the signs of regulation for the expression in plants.

- The two polypeptides are located in the endoplasmic reticulum and the protease 3ABC is capable of process the polypeptide Δ P1-2A and to achieve the expression of pentamers exclusively. These plants exhibit higher levels of expression as well as a better growth and development, which bears to obtain more biomass.

Identification of transgenic plants that express the product of the genes of the modified HAV.

A. tumefaciens was transformed with each of the binary vectors and there were obtained bacterial colonies containing these plasmids. *A. tumefaciens* independently carrying the

different genetic constructions were used for plant transformation and finally to obtain plants resistant to the kanamycin as a selection marker. The integration of the foreign DNA in plants was verified by the use of Southern blot and PCR techniques.

From leaves of transgenic plants, extraction of soluble proteins was performed, grinding them with liquid nitrogen in a particular protein extraction buffer. Capsids and pentamers were identified using specific antiserum to the HAV and a neutralizing monoclonal antibody. Immunological methods such as Western blot, ELISA or Immunomicroscopy were performed, they demonstrated that transgenic plants express the polyprotein or in some cases the expected polypeptides, and also that they are processed and assembled in pentamers or capsids.

Plants expressing higher levels of recombinant proteins were used for capsid and pentamer purification using a neutralizing monoclonal antibody.

Determination of the immunogenicity of the purified capsids and pentamers.

Capsid and pentamers immunopotency of HAV was determined by the immunological response of immunized mice with the purified product from plant leaves of tobacco and rice. Also the potency was evaluated in mice fed with transgenic carrots expressing the HAV antigen. Oral route and parenteral way were used as methods for the introduction of the antigen. The immune response was controlled and verified using Elisa's technology to determine the reactivity of the antiserum of the animal to the HAV, and for the neutralizing capacity of the immune sera to neutralize the HAV infection in vitro.

Advantages of the invention.

Among the most important advantages that our invention offers are: antigenic similarity among envelopes and purified pentamers as products of the expression of our constructs and the original virus; the expression levels of envelopes and pentamers in plants as products of the constructs we claim are higher either than the ones obtained when the HAV open reading frame is expressed or when the region P1-2A and the region P3 are co-expressed, because the polyprotein obtained as product of the expression of our constructs has a significantly lower size than the original virus size; the polypeptide 3ABC is exclusively composed by the proteins 3A, 3B and 3C and has mutated the selfprocessing sites between the proteins 3A/3B and 3B/3C, it avoids the polypeptide processing and therefore the proteolytic function of the polyprotein is assumed by the polypeptide itself with higher efficiency; pentamers and HAV viral envelopes expression levels in endoplasmic reticulum of the vegetable cell are higher than in the cytoplasm; the exclusive expression of pentamers in the vegetable cell is more efficient and allows a better plant growth and development due to the lower size of these

particles; the scale-up and production of pharmaceutical protein from plants is suitable to produce high amounts of antigens; production costs are reduced compared to other systems currently used and described in the state of the art; HAV antigen expression in plants decreases the contamination risks with pathogens affecting human beings; oral immunization against HAV significantly contributes to cheap the immunization costs due to the possibility to use plants without the need to purify the product.

Microorganism Deposit.

Plasmids pBVHARE, p Δ VHARE, pBMLAm and p Δ MLAm were leave to deposit under the regulations of Budapest Treatment for Microorganism Deposit at the Belgian Coordinated collection of Microorganisms BCCM, LMBP-COLLECTION with the access numbers LMBP 4721, LMBP 4722, LMBP 4723 and LMBP 4724 respectively deposited on May 19, 2003.

DESCRIPTION OF FIGURES.

Figure 1. Genetic constructs for envelopes and pentamers expression in plant vegetable cytosol. A) ORF scheme of the M2 strain of HAV. B) Scheme of the sequences that code for the structural proteins (P1-2A). C) Scheme of the sequences that code for the 3ABC region. D) Scheme of the ORFm of HAV. E) Scheme of the insert of interest cloned in the binary vector for plant expression.

Figure 2. Genetic constructs for pentamer expression in the cytosol of the vegetable cell. A) Scheme of the Δ ORFm without the sequence that codes for VP4. B) Scheme of the insert of interest cloned in the binary vector for Δ ORFm plant expression.

Figure 3. Genetic constructs for the expression of envelopes and pentamers in the endoplasmic reticulum of the vegetable cell. A) Scheme of the P1-2A sequence fused to KDEL sequence. B) Scheme of the insert of interest cloned in the binary vector for the expression in the endoplasmic reticulum. E- Spacer, K- KDEL

Figure 4. Genetic constructs for the expression of pentamers in the endoplasmic reticulum of the vegetable cell. A) Scheme of the P1-2A sequence without the VP4 sequence fused to KDEL sequence. B) Scheme of the insert of interest cloned in the binary vector for plant expression. E- Spacer, K- KDEL

Figure 5. Southern blot of genomic DNA from transgenic plants.

Figure 6. Southern blot of PCR products from carrot and rice transgenic plants.

Figure 7. Western blot of plant proteins from transgenic tobacco, carrot and rice transformed with genetic constructs for the expression in the cytosol of envelopes and pentamers.

Figure 8. Immune-enzymatic assay (ELISA) performed to tobacco, carrot and rice plants transformed with the genetic constructs for the expression in the cytosol of HAV envelopes and pentamers.

Figure 9. Electronic immune-microscopy of one tobacco plant, transformed with the construct pBMLAm. A) Untransformed plant. B) Transformed plant. C) Transformed plant.

Figure 10. Inhibition ELISA of the sera from mice immunized with HAV by intraperitoneal way.

Figure 11. Inhibition ELISA of the sera from mice orally immunized with HAV pentamers purified from rice and tobacco plants.

Figure 12. Inhibition ELISA of the sera from mice orally immunized by feeding them with carrots collected from plants expressing HAV pentamers.

EXAMPLES.

Example 1. Cloning of the ORF of the HAV of the Cuban strain M2.

The sequence of interest of the plasmid pMLA1 is shown in the Figure 1 (A). From the HAV strain M2 isolated and characterized in Cuba, the RNA was purified and a DNA fragment of 6.7 kb was amplified by means of Reverse Transcription-Polymerase Chain Reaction technology (RT - PCR), using specific oligonucleotides (SEQ ID NO 1 and 2) for other previously reported sequences of the HAV. The band was cloned in a vector BlueScript (KS+), previously digested with *Sma*I endonuclease. The resultant plasmid was named pMLA1 and was used for the sequencing determination of HAV's ORF of the Cuban M2 strain. DNA sequence (SEQ ID NO 3) presents changes with regard to the reported ones. It has changes that generate 11 different aminoacids. The analysis of these sequences allows to classify the strain M2 inside the subgenotype IA, to which belong almost the totality of the American strains. The sequence confirms that the Cuban strain M2 is a really HAV strain different from the ones previously reported.

Example 2. Genetic constructs for the expression of capsides and pentamers in the plant cell cytosol.

The sequence of interest of the plasmid pP1-2A is shown in the Figure 1 (B). This plasmid was obtained by PCR amplification of the sequences coding for the structural proteins (P1-2A), using specific oligonucleotides (SEQ ID NO 4) complementary for the 5' region of the sequence that codes for the protein VP4 and the 3' region of the sequence that codes for the protein 2A from the plasmid pMLA1, respectively. The amplified band of 2.5 kb (SEQ ID NO 6) was cloned in the vector BlueScript (KS+) *Sma*I digested.

To obtain the plasmid p3ABC, which sequence of interest is shown in the Figure 1 (C), the region of 0.2 kb that codes for the protein 3A was amplified by PCR, using oligonucleotides (SEQ ID NO 7 and 8) complementary for the 5' and 3' regions of this gene. It was cloned in a vector Blue Script (KS+) *Bam*HI/*Eco*RV digested. After that, in the same vector but in the region among *Eco*RV/*Xba*I sites, a synthetic nucleotide sequence (SEQ ID NO 9 and 10) coding for the protein 3B was cloned. The obtained plasmid was called p3AB. On the other hand, from the plasmid pMLA1, the sequence of 0.65 kb that codes for the protein 3C was amplified by PCR using the oligonucleotides SEQ ID NO 11 and 12. This band was cloned among the sites *Xba*I/*Hind*III of the vector P3AB. The resulting sequence obtained was called 3ABC (SEQ ID NO 13),

It codes for a polyprotein with proteolytic activity but without self-processing possibilities because the cutting sites between the proteins 3A/B and 3B/C are mutated by means of the nucleotide substitution T by C and G by C respectively.

The sequence of interest of the plasmid pMLAm is shown in the Figure 1 (D). To obtain it, the plasmid pP1-2A was *Eco*RI and *Cla*I digested. A 1kb band coding for the mutated polypeptide 3ABC was extracted by *Eco*RI/*Cla*I digestion and cloned in the proper sites of the pP1-2A previously digested. The plasmid pMLAm contains the modified sequence that codes for a HAV polyprotein of a significantly minor weight compared to the original polyprotein (SEQ ID NO 14).

The plasmid pKMLAm was obtained by cloning the 3.4 kb ORFm band *Sma*I/*Cla*I digested in the vector pKTPL-2. This vector contains as promoter 2X the sequence of 35SCaMV promoter, the leader sequence of the TEV and the terminator of the 35S CaMV. To clone the ORFm band, the plasmid pKTPL-2 was *Nco*I digested followed by blunting with *Klenow* fragment of DNA *Pol*I and finally *Cla*I digested.

The sequence of interest of the binary plasmid pBMLAm is shown in the Figure 1 (F). This plasmid was obtained by *Sph*I digestion of the plasmid pKMLAm and subsequent treatment with *Mung Bean* nuclease, rendering a band of 4.7 kb that was cloned in the binary vector pBin19 previously *Sma*I digested.

The resultant plasmid pBMLAm contains: the neomycin phosphotransferase II gene (NPTII) which acts as selection marker conferring kanamycin resistance; the ORFm gene which codes for the modified HAV polyprotein, regulated by the 2X 35S CaMV promoter and the leader of the TEV as well as the terminator of the CaMV. It also has the borders sequences of T-DNA to facilitate its transference to the plant genome.

Example 3. Genetic constructs for the expression of pentamers in the cytosol of the plant cell.

The sequence of interest of the plasmid p Δ MLAm is shown in the Figure 2 (A). To eliminate the protein VP4 and to obtain this plasmid, the fragment of 114bp was eliminated from the plasmid pMLAm cutting with the enzymes *SmaI/PstI* and it was replaced with the synthetic nucleotide sequence (SEQ ID NO 15 and 16) that returns the beginning of the gene that codes for the protein VP2. The sequence of the region Δ ORFm corresponds to the SEQ ID NO 17. The plasmid pK Δ MLAm was obtained cloning the band Δ ORFm (3.46 kb) *SmaI/ClaI* digested in the plasmid pKTPL-2 previously digested *NcoI/Klenow/ClaI*.

10 The sequence of interest of the binary plasmid pB Δ MLAm is shown in the Figure 2 (B), the binary plasmid was obtained by digestion of Bin19 with the enzyme *SmaI* and a DNA fragment of 4.6 kb resultant of the digestion of the plasmid pKMLAm with the enzyme *SphI* and treatment with *Mung Bean* nuclease was then cloned.

The resultant plasmid pB Δ MLAm contains: the neomycin phosphotransferase II (NPT II) gene which functions as selectable marker conferring kanamycine resistance; the ORFm gene which codes for the HAV modified polyprotein, regulated by the sequence 2X 35S CaMV as promoter and TEV leader, as well as the terminator of 35S CaMV. It also contains the right and left borders to be transferred to the plant genome.

20 **Example 4. Genetic constructs for the expression of capsides and pentamers in the endoplasmic reticulum of the plant cell.**

The sequence of interest of the plasmid pBVHARE appears in the Figure 3B. To obtain this plasmid, a synthetic fragment (SEQ ID NO 18 and 19) that codes for KDEL retention signal was cloned in the sites *EcoRV/ClaI* of the vector BS (+). On the other hand, in the sites *StyI/EcoRI* of the plasmid pP1-2A, a synthetic fragment (SEQ ID NO 20 and 21) that modifies the 3' end of the protein 2A and eliminates the protease cutting site in this region, as well as introduces a sequence that works as space-bar between the union of the gene and the sequence that codes for the KDEL signal, was cloned. Later this sequence (2.5 kb) was extracted with the enzymes *SmaI/EcoRV* and was cloned in the vector BS-KDEL, resulting the plasmid pP1-2ARE (Figure 3A, SEQ ID NO 22). The plasmid p3ABCRE was obtained digesting the p3ABC plasmid with the enzymes *XhoI/Klenow/EcoRI* and cloning the 3ABC sequences in the *EcoRI/EcoRV* sites (Figure 3A, SEQ ID NO 23).

30 To provide plant expression regulatory signals to the genes of interest, the structural region P1-2A-KDEL (2.5 kb), extracted from the plasmid pP1-2ARE with the enzymes *SmaI/ClaI*, was cloned in the plasmid pKTPL-2 digested *NcoI/Klenow/ClaI*. The resulting plasmid was

named pKP1-2ARE. The 1kb region 3ABC-KDEL was extracted from the plasmid p3ABCRE with the enzymes *NcoI/ClaI* and was cloned in the plasmid pKTPL-2 digested with the same enzymes. The resulting plasmid was pK3ABCRE.

Finally, to achieve the plasmid for plant transformation by means of *A. tumefaciens*, the 2 kb
 5 sequence extracted from the plasmid pK3ABCRE with enzyme *Sall* was cloned in the binary vector pBin 19 previously digested with the same enzyme. Resulting plasmid was called pB3ABCRE. Later, in the *SphI* site of the same vector, the expression cassette corresponding to the sequence P1-2A-KDEL, extracted from the plasmid pKP1-2ARE by *SphI* digestion, was cloned. The resultant plasmid pBVHARE carries separated both the structural regions
 10 and the region with protease function, fused to the retention signals in the reticulum under plant expression regulatory signals and the neomycin phosphotransferase II (NPT II) gene as selectable marker.

Example 5. Genetic constructs for the expression of pentamers in the endoplasmic reticulum of the plant cell.

15 The sequence of interest of the plasmid p Δ VHARE is shown in the Figure 4(B). This plasmid was obtained cutting the plasmid pP1-2ARE with the enzymes *SmaI/PstI* and it was replaced with the synthetic nucleotide sequence (SEQ ID NO 15 and 16) that returns the beginning of the gene that codes for the protein VP2. The resultant plasmid p Δ P1-2ARE (Figure 4A, SEQ ID NO 24), was digested *SmaI/ClaI* and the band of the 2.4 kb was cloned
 20 in the vector pKTPL-2 digested *NcoI/Klenow/ClaI*, resulting the plasmid pK Δ P1-2ARE. It was cloned in the plasmid pB3ABCRE (the binary plasmid containing the 3ABC-KDEL), the expression cassette digested with the *SphI* enzyme).

The resultant binary plasmid contains: the structural region, without the sequences that code for the protein VP4, fused to the KDEL peptide, under plant expression signals; the 3ABC-KDEL region under the same signals and the neomycin phosphotransferase II (NPT II) gene
 25 as the selectable marker.

Example 6. Obtention of capsids and pentamers of HAV in *Nicotina tabacum* transgenic plants.

The genetic transformation of *Nicotiana tabacum* plants was carried out following the
 30 method of Zambryski *et al.*, (1983).

The *A.tumefaciens* strain At 2260 (Deblaere *et al.*, 1985) was transformed by the method of the liquid nitrogen (Hofgen and Willmitzer, 1988) with the developed binary plasmids (P Δ VHARE , PBVHARE, P Δ MLAm, PBMLAm). Leave discs of tobacco plants of the

variety Petit Havana MR 1 cultivated “in vitro” were transformed with the recombinant *Agrobacteriums*. Kanamycine (100 mg/L) was used as selection marker.

Several procedures such as Southern blot, Western blot, ELISA and immunomicroscopy were performed to detect the presence of the genes of interest in the tobacco plant genome and their expression, as well as formation of either viral envelop or pentamers.

Example 7. Obtention of capsides and pentamers of HAV in transgenic carrot plants (*Daucus carota* L.)

For the transformation of this plant were used The *A. tumefaciens* strain At 2260 transformed with the binary plasmids (PBΔVHARE , PBVHARE, PBΔMLAm, PBMLAm) was used for plant transformation. Germinated three weeks old hypocotyls of the variety NEW KURODA were cut in 1 cm segments and planted in BAN-9 medium (Murashige and Skoog, 1962 (MS), supplemented with 0.5 mg/L of NAA) for three days. Later they were incubated during 30 minutes with an At suspension containing each one of the constructions previously described. Again, explants were transferred to BAN-9 medium for 72 hours. After this period, they were planted in regeneration medium supplemented with Kanamycine (100 mg/l). Shoots appeared after 3 weeks and they were individualized and planted in MS medium also supplemented 100 mg/L of Kanamycine. Gene integration was verify by Southern blot of PCR products (Figure 6). Polyprotein expression, its processing and formation of viral capsids and pentamers was demonstrated by ELISA (Figure 8) and Western blot (Figure 7).

Example 8. Obtaining of the capsid and pentamers of the HAV in transgenic rice plants (*Oryza sativa* L.).

The genetic transformation of rice plants was carried out following the method used by Hiei *et al.*, (1994). The *A.tumefaciens* strain At2260 was transformed with the developed binary plasmids (pBΔVHARE, pBVHARE, pBΔMLAm, pBMLAm) using method of the liquid nitrogen. Callus obtained from rice escutelo were transformed with recombinant *A. tumefaciens*. Kanamycine (100 mg/L) was used as selection marker.

To verify the presence of the genes of interest in plant rice genome and their expression, as well as the formation of viral capsids or pentamers, different procedures were performed, which are below described.

Example 9. Molecular characterization and immunochemical of the transgenic plants.

Analysis by Southern blot.

The method described by Dellaporta *et al.*, (1983) was used to obtain chromosomal DNA for Southern blot purposes from tobacco, carrot and rice plants. As samples, leaves of

transformed plants with the previously described constructs that showed resistance to the selection marker were used. Leaves of untransformed plants were used as negative control.

Total DNA digestions, agarose gel electrophoresis, transfer to a Hybond N membrane and hybridization were performed by standard procedures (Sambrook *et al.*, 1989). A 1.2 kb

5 DNA fragment that includes the gene coding for VP1 protein VP1 was ^{32}P labeled by means of a Primer-a-Gene Labeling System (Promega Corp., the USA) and used as radioactive probe. The same fragment was used as positive control.

Figure 5 shows the Southern blot of transgenic tobacco plants transformed with the constructs PBΔMLAm and PBMLAm to expressing in the cytoplasm both capsids and pentamers of the
10 HAV. By DNA digestion with *Sma*I and *Cl*aI there is a resulting band of 3.4 kb. Total DNA from transgenic plants transformed with the constructs PBΔVHARE and PBVHARE to be expressed in the endoplasmic reticulum was digested with *Sma*I I-*Eco*RI resulting a band of 2.4 kb. Results shown in the Figure 5 demonstrate that plants contain in their genomes the sequences coding for the structural proteins.

15 Both in transgenic carrot and rice plants, a Southern blot was performed to the PCR amplification products with oligonucleotides corresponding to the sequences SEQ ID NO 4 and 5. As it is shown in the Figure 6, the radioactively labeled sequence that codes for the protein VP1 complements with a band of 2.5 kb corresponding to the expected size of the sequence that codes for the structural proteins.

20 Analysis by Western blot.

Western blot results are shown in the Figure 7. Western blot assay for immunodetection of the recombinant molecules was carried out according to the methodology described by Towbin *et al.*, (1979). Western blot samples consisted on total soluble proteins extracted from transgenic tobacco, carrot and rice plants, transformed with the constructs to express only
25 pentamers: Clones tobacco 5, carrot 7 and rice 3, transformed with the construct pBΔVHARE to express pentamers in the endoplasmic reticulum; and clones tobacco 25 and carrot 10, transformed with the construct pBΔMLAm which allows the expression of the pentamers in the cytoplasm. As negative controls, protein extracted from untransformed tobacco leaves were used. As positive control, the VP1 protein expressed in *E.coli*. Leaves were grinded
30 with liquid nitrogen up to obtain a very thin dust. 1 mL of protein extraction buffer [Tris-HCl 61 mM pH 6.8, Triton 0.1%, glycerol 12.5% and Fluoride of Fenilmetilsulfonilo (PMSF) 1 mM] per gram of leaves was added as reported Schouten *et al.*, (1997). The insoluble material was removed by centrifugation at 13 000 rpm.

Total proteins of SDS-PAGE were transferred to a nitrocellulose membrane and proteins of interest were identified using a polyclonal antibody anti-VP1 conjugated to the enzyme alkaline phosphatase (PhoA). The enzyme detection was performed by means of a colorimetric reaction.

- 5 In the Figure 7 is possible to observe the presence of a protein band of the same size of the VP1 protein in all cultures, as well as other intermediate products resulting from incomplete processing of the polyprotein.

Immunoenzymatic test (ELISA).

- ELISA results are shown in the Figure 8. "Sandwich" assays were performed. The plate
10 (Maxisorp, Nunc) was covered with 10 mg/mL of the monoclonal antibody 7E7 in carbonate buffer (Na_2CO_3 0.015 M, NaHCO_3 0.028 M, pH 9.6) during 4 hours at 37 °C. Blocking was carried out during 2 hours at 37 °C, with 5% of milk in PBS (NaCl 100 mM, Na_2PO_4 80 mM, NaH_2PO_4 20 mM, pH 7.4). Later 100 μL of the samples corresponding to transformed and untransformed tobacco, carrot and rice plants (prepared in the same way as was described for
15 the Western blot), were added. Plate was incubated overnight at 4°C. After washing with PBS, 100 μL of the monoclonal antibody 7E7, conjugated with alkaline phosphatase at 1/1000 dilution (1 mg/mL in PBS containing 0.5% milk) were added. The plate was incubated at 37 °C during 1 hour. The reaction was developed by addition of 4-nitrophenylphosphate (substrate of the enzyme), prepared in 0.1% of Diethanolamine. Color
20 appearance was followed during a 60 minutes period. The absorbance was read at a wavelength of 405nm in a spectrophotometer. The washes of the plate in every stage of the ELISA were performed three times with PBS containing 0.1% of Tween 20.

Analysis for electronic immunomicroscopy.

- Immunomicroscopy results are shown in the Figure 9. Samples of transformed tobacco with
25 the plasmid pBMLAm and untransformed plants, both from tissue culture, were fixed in a solution of 4% of Formaldehyde and later in 0.2% of Glutaraldehyde. They were dehydrated in ethanol and then incubated in a Lowicryl K 4M (Chemische Werke Lowi, Waldkraiburg) solution. Ultra thin cuts were placed in a nickel grill and incubated with the monoclonal antibody 7E7. After this step, there was an incubation period with the polyclonal antibody
30 anti-mouse IgG labeled with 15 nm of gold colloidal particles (British Bio-Cell International). Immunolabeled sections were contrasted by a 5 minutes treatment in Uranilacetate followed by 7 minutes in Lead Citrate, before being examined by a transmission electronic microscope (Jeol-Jem 2000EX, Japan). Results show particles of

approximately 27 nm of diameter only in the tobacco transformed plants with the construction pBMLAm, by means of what the protein was expressed in the cell cytoplasm.

Example 10. Purification of capsids and pentamers from transgenic tobacco and rice plants.

- 5 For capsid and pentamer purification a monoclonal antibody anti HAV obtained at CIGB laboratories that exclusively recognizes the particles and immunogenic pentamers was used. Vegetable cell proteins were extracted using the protocol described for Western blot analysis. Supernatant resultant from centrifugation was dissolved in 0.5M of sodium chloride and mixed with an affinity gel (Bio-rad Laboratories, Richmond, CA) charged with an antibody.
- 10 The mix was incubated for 16 hours at 4°C. The gel was washed with 10 volumes of PBS (NaCl 100 mM, Na₂PO₄ 80 mM, NaH₂PO₄ 20 mM, pH 7.4) and later the protein of interest was eluted with glycine 0.2 M at pH 2.5. The eluate was neutralized with basic Tris and was dialyzed against PBS. The presence of HAV particles and pentamers from these leave extracts was detected by ELISA, using a 7E7 commercial monoclonal antibody
- 15 (Mediagnost), specific for recognition of HAV viral capsids and pentamers.

Example 11. Determination of the immunogenicity of the capsids and pentamers purified from transgenic plants by means of the intraperitoneal administration.

- White ICR mice of 14 weeks old were immunized with two doses of 750 EL.U of capsids and pentamers purified from transgenic tobacco and rice plants. In the same way, a group of mice
- 20 were inoculated with a commercial HAV antigen (Mediagnost) and used as positive control. Another group was inoculated with PBS and used as negative control. Blood samples were collected 0, 15, 30, 50 and 70 days post-inoculation.

- The antibody levels were measured by means of an inhibition ELISA: the plate was covered with 5 µg of the monoclonal antibody 7E7 and then incubated for 4 hours. Later it was
- 25 washed once with PBS-0.1% Tween. Blocking was performed adding 5 % of milk in PBS-0.1% Tween and incubated for 2 hours at 37 °C. The plate was washed 3 times with PBS-0.1% Tween. Sera from immunized mice, previously incubated during 20 min at 37°C, with HAV antigen (Mediagnost), were added. The plate was incubated for 12 hours at 16°C and washed 5 times with PBS-0.1% Tween. Finally, 100µL of the monoclonal antibody 7E7
- 30 conjugated with alkaline phosphatase diluted 1/1000 in PBS containing 0.5% milk, were added. The incubation was carried out during an hour at 37°C. The reaction was developed by addition of 4-nitrophenylphosphate (enzyme substrate) prepared in diethanolamine. Color appearance was followed during 60 minutes. The absorbance was read to a wavelength of 405 nm in an spectrophotometer. In the Figure 10 is shown the average levels of inhibition of

the sera of the mice inoculated with the antigen purified from transgenic plants, detected in blood of the mice immunized with pentamers produced by tobacco and rice plants. In the same way, similar levels of antibody were observed in mice immunized with the antigen produced in tobacco and rice plants transformed with the constructs that allow the expression of capsids and pentamers.

Example 12. Determination of the immunogenicity of the capsids and pentamers purified from transgenic plants by means of the oral administration.

The antigen oral administration was performed by two routes: using the purified antigen and feeding animals with carrots expressing the antigen.

- For antigenicity determination of purified capsids and pentamers administered by oral way, pentamers and capsids were administered to 8 weeks Balb/c mice in four doses of 7500 EL.U. 200 μ L of blood at 0, 15, 30, 50 and 70 days post - inoculation were collected to detect the presence of anti HAV antibodies by an ELISA of inhibition.

- ELISA of inhibition was performed by means of the procedure described previously in the example 11.

- According to the results shown in the Figure 11, the oral administration of the HAV pentamers expressed in transgenic plants produces immune response that is demonstrated by average inhibition of the sera from mice used in the experiment. Average inhibition of the sera from mice orally administered were lower compared with the ones obtained after intraperitoneal administration. Pentamers oral administration through plants was carried out with 5 g of row carrots (transformed with the construct pB Δ MLAm specifically designed to produce only pentamers) once a week during 4 weeks., sera from mice fed with untransformed carrots, were used as negative control. Ability of these plants to rise an immune response was demonstrated by an inhibition ELISA shown in the Figure 12.

SEQUENCE LISTING

5 <110> CENTER FOR GENETIC ENGINEERING AND BIOTECHNOLOGY
 <120> RECOMBINANT HEPATITIS A VIRUS ANTIGENS PRODUCED IN PLANT CELLS.

 10 <130> ORF.
 <140>
 <141>

 15 <160> 24
 <170> PatentIn Ver. 2.1

 20 <210> 1
 <211> 25
 <212> ADN
 <213> Chimeric Sequence

 25 <220>
 <221> primer_bind
 <222> (1)..(25)
 <223> Sequence # 1.
 Sequence of the oligonucleotide # 1 used for the amplification of ORF
 coding sequence by RT-PCR.
 30

 <400> 1
 cttaatctag aatgaatatg tccaa 25

 35 <210> 2
 <211> 22
 <212> ADN
 <213> Chimeric Sequence

 40 <220>
 <221> primer_bind
 <222> (1)..(22)
 <223> Sequence # 2.
 45 Sequence of the oligonucleotide # 2 used for the amplification of ORF
 coding sequence by RT-PCR.
 <400> 2
 gaaagaaata aaggtacctc ag 22

 50 <210> 3
 <211> 6685
 <212> ADN
 <213> Hepatitis A virus

 55 <220>
 <221> gene
 <222> Complement((1)..(6685))
 <223> Sequence # 3.
 60 Nucleotide sequence coding for the HAV open reading frame (ORF) of
 the Cuban M2 strain.

<400> 3

	atgaatatgt	ccaaacaagg	aattttccag	actggtggga	gtggccttga	ccacatcctg	60
5	tccttggcag	atattgagga	agagcaaagt	attcagtcg	ttgataggac	tgcagtgact	120
	ggagcttctt	atttcacttc	tgtggaccaa	tcttcagttc	atactgctga	ggttggctca	180
	caccaaattg	aacctttgaa	aacctctgtt	gataaacctg	gttctaagaa	aactcagggg	240
	gagaagtttt	tcttgattca	ttctgctgat	tggctcacta	cacatgctct	ctttcatgaa	300
	gttgcaaaat	tggatgtggg	gaaactgctg	tacaatgagc	agtttgccgt	ccaaggtttg	360
10	ttgagatacc	atacttatgc	aagatttggc	attgagattc	aagttcagat	aaatcccaca	420
	ccctttcagc	aaggaggact	aatctgtgcc	atggttcctg	gtgaccaaag	ttatggttca	480
	atagcatcct	tgactgttta	tcctcatggg	ctgttaaatt	gcaatatcaa	caatgtagtt	540
	agaataaagg	ttccatttat	ttatactaga	ggtgcttata	attttaaaga	tccacagtac	600
	ccagtttggg	aattgacaat	cagagtttgg	tcagagttga	atattggaac	aggaacctca	660
15	gcttatactt	cactcaatgt	tttagctagg	tttacagatt	tggagttgca	tggattaact	720
	cctctttcta	cacagatgat	gagaaatgaa	tttagagtta	gtactactga	aaatgttgta	780
	aatttgtcaa	attatgaaga	tgcaagggca	aaaatgtctt	ttgctttgga	tcaggaagat	840
	tggaagtctg	atccttccca	aggtggtgga	attaaaatta	ctcatttcac	tacctggaca	900
	tccattccaa	ccttagctgc	tcagtttcca	ttcaatgctt	cagattcagt	tgggcaacaa	960
20	attaaagtta	taccagtgga	cccatacttt	ttccagatga	caaacactaa	tcctgatcaa	1020
	aaatgtataa	cagccttggc	ctctattttc	cagatgttct	gcttttggag	gggagatctt	1080
	gttttcgatt	tccagggttt	tccaaccaaa	tatcattcag	gtaggctgtt	gtttgttttt	1140
	gttcctggga	atgagttaat	agatgttact	ggaattacat	taaaacaggc	aactactgct	1200
	ccttgtgcag	tgatggacat	tacaggagtg	cagtcaacct	tgagatttcg	tgttccttgg	1260
25	atctctgata	cacctatcg	agtgaatagg	tacacgaagt	cagcacatca	aaaaggtgag	1320
	tatactgcc	ttgggaagct	tattgtgtat	tggtataata	gattgacttc	tccttctaata	1380
	gttgcttctc	atgttagagt	taatgtttat	ctttcagcaa	ttaattttgga	atgttttgct	1440
	cctctttacc	atgctatgga	tgttaccaca	caggttggag	atgattcagg	aggttttctca	1500
	acaacagttt	ctacagagca	gaatgttctc	gatccccaag	ttggcataac	aaccatgagg	1560
30	gatttaaaag	ggaaaagcaa	taggggaaag	attgatgtat	caggagtgca	ggtacctgtg	1620
	ggagctatta	gacaatgga	ggatccagtt	ttagcaaaaga	aagtacctga	gacatttcct	1680
	gaattgaagc	ctggagaatc	cagacataca	tcagatcaca	tgtctattta	taaattcatg	1740
	ggaaggtctc	atttcttgtg	tacttttact	tttaattcaa	acaataaaga	gtacacattt	1800
	ccaataactc	tgtcttcgac	ttctaatacct	cctcatgggt	taccatcaac	attaaggtgg	1860
35	ttctttaatt	tgtttcagtt	gtatagagga	ccattggatt	tgacaattat	aatcacagga	1920
	gccactgatg	tggatgggat	ggcctgggtt	actccagtg	gccttgctgt	cgacacccct	1980
	tgggtggaaa	agaagtcagc	tttgtctatt	gattataaaa	ctgcccttgg	agctgttaga	2040
	tttaatacaa	gaagaacagg	gaacattcag	attagattgc	catgggtattc	ttatttgtat	2100
	gccgtgtctg	gagcactgga	tggcttggga	gataagacag	attctacatt	tggattgggt	2160
40	tctattcaga	ttgcaaatga	caatcattct	gatgaatatt	tgtcctttag	ttgttatttg	2220
	tctgtcacag	agcaatcaga	gttctatttc	cctagagctc	cattaaattc	aaatgctatg	2280
	ttgtccactg	agtccatgat	gagtagaatt	gcagctggag	acttgagtc	atcagtggat	2340
	gatcccagat	cagaggagga	cagaagattt	gagagtcata	tagaatgtag	gaaaccatat	2400
	aaagaattga	gactggagg	tgggaaacaa	agaatcaaat	atgctcagga	agagttatca	2460
45	aatgaagtgc	ttccacctcc	taggaaaatg	aaggggttat	tttcacaagc	taaaatttct	2520
	cttttttata	cagaggacca	tgaataatg	aaattttctt	ggagaggagt	gactgctaata	2580
	actagggtt	tgagaagatt	tggattctct	ctggctgctg	gtagaagtgt	gtggactctt	2640
	gaaatggatg	ctggagttct	tactggaaga	ttgatcagat	tgaatgatga	gaaatggaca	2700
	gaaatgaagg	atgataagat	tgtttcatta	attgaaaagt	tcacaagcaa	taaacattgg	2760
50	tctaaagtga	attttccaca	tggaaatgtg	gatcttgagg	aaattgctgc	caactctaaa	2820
	gatttttcaa	atatgtctga	gacagatttg	tgtttctgt	tgcattggct	aaatccaaag	2880
	aaaattaatt	tagcagatag	aatgcttggg	ttgtctggag	tgcaggaaat	taaagaacag	2940
	ggtgttggac	tgatagcaga	gtgtagaact	ttcttggatt	ctattgctgg	gactttgaaa	3000
	tctatgattt	ttgggtttca	ttattctgtg	actgttgaaa	ttataaatat	tgtgctttgt	3060
55	tttattaaga	gtggaatcct	gctttatgtc	atacaacaat	tgaaccaaga	tgaacactct	3120
	cacataattg	gtttgttgag	agttatgaat	tatgcagata	ttggctgttc	agtcatttca	3180
	tgtggtaaag	ttttttccaa	aatgttagaa	acagttttta	attggcaaat	ggactctaga	3240
	atgatggagc	tgaggactca	gagcttctcc	aattggttaa	gagataattg	ttcggaattt	3300
	actattttta	aaagttttta	ggatgccata	tattggttat	gtacaaaatt	gaaggatttt	3360
60	tatgaagtaa	attatggcaa	gaaaaaggat	gttcttaata	ttctcaaaga	taaccagcaa	3420
	aaaatagaaa	aagccattga	agaagcagac	aatttttgca	ttttgcaaat	tcaagatgtg	3480
	gagaaatttg	atcagtatca	gaaaggggtt	gatttaatac	aaaagctgag	aactgtccat	3540

5	tcaatggctc	aagttgaccc	cagtttgggg	gttcatttgt	cacctctcag	agattgcata	3600
	gcaagagtcc	atcaaaaagct	caagaatctt	ggatctataa	atcaggccat	ggtaacaaga	3660
	tgtgagccag	ttgtttgcta	tttgtatggc	aaaagagggg	gagggaaaag	cttgacttca	3720
	attgcattgg	caaccaaagt	ttgtaaacac	tatggtgttg	aacctgagaa	aaatatattac	3780
	accaaacctg	tggcctcaga	ttattgggat	ggatatagtg	gacaattagt	ttgtattatt	3840
10	gatgatatcg	gccaaaacac	aacagatgaa	gattggtcag	atttttgtca	attagtggtca	3900
	ggatgcccaa	tgagattgaa	tatggcttct	cttgaggaga	agggcagaca	tttttctctt	3960
	ccttttataa	tagcatcttc	aaattgggtca	aatccaagtc	caaaaacagt	ttatgttaaa	4020
	gaagcaattg	atcgtaggct	tcatttttaag	gttgaagtta	aacctgcttc	atttttttaa	4080
	aatcctcaca	atgatattgt	aaatgttaat	ttggctaaaa	caaattgatgc	aattaaagac	4140
15	atgtcttgtg	ttgatttgat	aatggatgga	cacaatat	cattgatgga	tttacttagt	4200
	tccttagtga	tgacaggtga	aattaggaaa	cagaatatga	gtgaattcat	ggagttgtgg	4260
	tctcagggaa	tttcagatga	tgacaatgat	agtgcagtag	ctgagttttt	ccggtctttt	4320
	ccatctgggtg	aaccatcaaa	ttccaagtta	tctagttttt	tccaagctgt	cactaatcac	4380
	aagtgggttg	ctgtggggagc	tgcagttggt	attcttggat	tgctagtggg	aggatgggtt	4440
20	gtgtataagc	atttttcccg	caaagaggaa	gaaccaattc	cagctgaagg	ggtttatcat	4500
	ggagtgcata	agcccaaaca	agtgtattaa	ttggatgcag	atccagtaga	gtctcagtc	4560
	actctagaaa	tagcaggatt	agttaggaaa	aatttgggtc	agtttggagt	tggtgagaaa	4620
	aatggatgtg	tgagatgggt	catgaatgcc	ttaggagtga	aggatgattg	gttgttagta	4680
	ccttctcatg	cttataaaat	tgaaaaggat	tatgaaatga	tggagtttta	tttcaataga	4740
25	ggtggaactt	actattcaat	ttcagctggg	aatgttggtt	ttcaatcttt	agatgtggga	4800
	ttccaagatg	ttgttcta	gaaggttcct	acaattccca	agtttagaga	tattactcaa	4860
	cattttatta	agaaaggaga	tgtgcctaga	gccttgaatc	gcttggcaac	attagtgaca	4920
	accgttaatg	gaactcctat	gttaatttct	gagggacott	taaaaatgga	agaaaaagcc	4980
	acttatgttc	ataagaagaa	tgatggtact	acggttgatt	tgactgtaga	tcaggcatgg	5040
30	agaggaanaag	gtgaaggtct	tcctggaatg	tgtggtgggg	ccctagtgtc	atcaaatcag	5100
	tccatacaaaa	atgcaatttt	gggtattcat	gttgctggag	gaaattcaat	tcttgtggca	5160
	aagttgatta	ctcaagaaat	gtttcaaaac	attgataaga	aaattgaaag	tcagagaata	5220
	atgaaagtgg	aatttactca	atgttcaatg	aatgtagtct	ccaaaacgct	tttttagaaag	5280
	agtcccatct	atcaccacat	tgataaaacc	atgattaatt	ttcctgcagc	tatgcctttc	5340
35	tctaaagctg	aaattgatcc	aatggctatg	atgttgctta	aatattcatt	acctattgtg	5400
	gaagaaccag	aggattacaa	agaagcttca	gttttttatt	aaaataaaat	agtaggcaag	5460
	actcagctag	ttgatgactt	tctagatctt	gatattggcca	ttacaggggc	tccaggcatt	5520
	gatgctatta	atatggattc	atctcctggg	tttccttatg	ttcaagaaaa	attgactaaa	5580
	agagatttga	tttggttgga	tgaaaatggt	ttaactgttag	gagttcacc	aagattggcc	5640
40	cagagaatct	tattttaatac	tgtcatgatg	gaaaattggt	ctgacttaga	tgttgttttt	5700
	acaacttgct	caaaaagatga	attgagacca	ttagagaaaag	ttttggaatc	aaaaacaaga	5760
	gctattgatg	cttgcccttt	ggattatata	attttatgtc	gaatgtattg	gggtccagct	5820
	attagttatt	ttcatttgaa	tccagggttt	cacacaggtg	ttgctattgg	catagatcct	5880
	gatagacagt	gggatgaatt	atttaaaaca	atgataagat	ttggagatgt	tggtcttgat	5940
45	ttagattttt	ctgctttttga	tgccagctct	agtccattta	tgattaggga	agcaggtaga	6000
	atcatgagtg	aattatctgg	aacaccatct	catttttgga	cagctcttat	caatactatc	6060
	atattattcta	aacatctgct	gtacaattgt	tgttatcacg	tctgtgggtc	aatgccttct	6120
	gggtctcctt	gtacagcttt	gttgaattca	attattaata	atattaattt	gtattattgtg	6180
	ttttctaaaa	tatttgga	gtctccagtt	ttctttttgtc	aagcttttag	gatcctttgt	6240
50	tatggagatg	atgttttgat	agttttttcc	agagatgttc	aaattgataa	tcttgacttg	6300
	attggacaga	aaattgtgga	tgagttcaaa	aaacttgga	tgacagccac	ttcagctgac	6360
	aaaaatgtgc	ctcaactgaa	gccagtttca	gaattgactt	ttcttaaaag	atctttta	6420
	ttggtggagg	acagaatcag	acctgcaatt	tcagaaaaga	caatttggtc	tttgatagct	6480
	tggcagagaa	gtaacgctga	gtttg				

<222> (1)..(40)

<223> Sequence # 4.

Sequence of the oligonucleotide # 5 used for the amplification of P1-2A coding sequence by PCR.

5

<400> 4

ttgaattcag cttgtgaaaa taacccttc attttcctag

40

10

<210> 5.

<211> 28

<212> ADN

<213> Chimeric Sequence

15

<220>

<221> primer_bind

<222> (1)..(28)

<223> Sequence # 5.

20

Sequence of the oligonucleotide # 5 used for the amplification of P1-2A coding sequence by PCR.

<400> 5

25

cgcccgggtc tagaatgaat atgtccaa

28

<210> 6

<211> 2523

30

<212> ADN

<213> Hepatitis A virus

<220>

<221> gene

35

<222> Complement((1)..(2523))

<223> Sequence # 6.

Nucleotide sequence coding for the structural P1-2A HAV proteins of the M2 strain.

40

<400> 6

atgaatatgt ccaaacaagg aattttccag actgttggga gtggccttga ccacatcctg 60
 tccttggcag atattgagga agagcaaag attcagtcg ttgataggac tgcagtgact 120
 ggagcttctt atttcacttc tgtggaccaa tcttcagttc atactgctga ggttggctca 180
 45 caccaaattg aacctttgaa aacctctgtt gataaacctg gttctaagaa aactcagggg 240
 gagaagtttt tcttgattca ttctgctgat tggctcacta cacatgctct ctttcatgaa 300
 gttgcaaaat tggatgtggt gaaactgctg tacaatgagc agtttgccgt ccaagggttg 360
 ttgagatacc atacttatgc aagatttggc attgagattc aagttcagat aaatcccaca 420
 ccctttcagc aaggaggact aatctgtgcc atggttccctg gtgaccaaag ttatggttca 480
 50 atagcatcct tgactgttta tcctcatggt ctgttaaatt gcaatatcaa caatgtagtt 540
 agaataaagg ttccatttat ttatactaga ggtgcttatt attttaaaga tccacagtac 600
 ccagtttggg aattgacaat cagagtttgg tcagagttga atattggaac aggaacctca 660
 gcttatactt cactcaatgt tttagctagg tttacagatt tggagttgca tggattaact 720
 cctctttcta cacagatgat gagaaatgaa tttagagtta gtactactga aaatgttgta 780
 55 aatttgtcaa attatgaaga tgcaagggca aaaatgtctt ttgctttgga tcaggaagat 840
 tggaagtctg atccttccca aggtggtgga attaaaatta ctcatctcac tacctggaca 900
 tccattccaa ccttagctgc tcagtttcca ttcaatgctt cagattcagt tgggcaacaa 960
 attaaagtta taccagtgga ccctactttt ttccagatga caaacactaa tcctgatcaa 1020
 aaatgtataa cagccttggc ctctatttgt cagatgttct gcttttggag gggagatctt 1080
 60 gttttcgatt tccaggtttt tccaacaaa tatcattcag gtaggctgtt gttttgtttt 1140
 gttcctggga atgagttaat agatgttact ggaattacat taaaacaggc aactactgct 1200
 ccttgtgcag tgatggacat tacaggagtg cagtcaacct tgagatttcg tgttccttgg 1260

```

atttctgata caccctatcg agtgaatagg tacacgaagt cagcacatca aaaaggtgag 1320
tatactgcc a ttgggaagct tattgtgtat tggtataata gattgacttc tccttctaata 1380
gttgcttctc atggttagagt taatgtttat ctttcagcaa ttaatttga atgttttctc 1440
cctctttacc atgctatgga tgttaccaca cagggtggag atgattcagg aggttttctc 1500
5 acaacagttt ctacagagca gaatgttcct gatccccaag ttggcataac aacctgagg 1560
gatttaaaag ggaaagccaa taggggaaag atggatgtat caggagtgc ggtacctgtg 1620
ggagctatta caacaattga ggatccagtt ttagcaaaga aagtacctga gacatttcct 1680
gaattgaagc ctggagaatc cagacatata tcagatcaca tgtctattta taaattcatg 1740
ggaagggtctc atttcttgtg tacttttact ttaattcaa acaataaaga gtacacattt 1800
10 ccaataactc tgtcttcgac ttctaatacct cctcatgggt taccatcaac attaggtgg 1860
ttctttaatt tggttcagtt gtatagagga ccattggatt tgacaattat aatcacagga 1920
gocactgatg tggatgggat ggcctgggtt actccagtgg gccttgctgt cgacaccct 1980
tggggtggaaa agaagtcagc tttgtctatt gattataaaa ctgcccttgg agctgttaga 2040
tttaatacaa gaagaacagg gaacattcag attagattgc catggtattc ttatttgtat 2100
15 gccgtgtctg gagcactgga tggcttggga gataagacag attctacatt tggattgggt 2160
tctattcaga ttgcaaatta caatcattct gatgaatatt tgccttttag ttgttatttg 2220
tctgtcacag agcaatcaga gttctatttc cctagagctc cattaaattc aaatgctatg 2280
ttgtccactg agtccatgat gagtagaatt gcagctggag acttgagtc atcagtggat 2340
gatcccagat cagaggagga cagaagattt gagagtcata tagaatgtag gaaaccatat 2400
20 aaagaattga gactggaggt tgggaaacaa agaatcaaat atgctcagga agagttatca 2460
aatgaagtgc ttccacctcc taggaaaatg aaggggttat atgcttctgg aggtgaattc 2520
gat 2523

```

```

25 <210> 7
    <211> 27
    <212> ADN
    <213> Chimeric Sequence

```

```

30 <220>
    <221> primer_bind
    <222> (1)..(27)
    <223> Sequence # 7.
        Sequence of the oligonucleotide # 7 used for
35 the amplification of 3A coding sequence by PCR.

```

```

    <400> 7
40 ccatgggaat ttcagatgat gacaatg 27

```

```

    <210> 8
    <211> 26
    <212> ADN
45 <213> Chimeric Sequence

```

```

    <220>
    <221> primer_bind
    <222> (1)..(26)
50 <223> Sequence # 8.
        Sequence of the oligonucleotide # 7 used for
the amplification of 3A coding sequence by PCR.

```

```

55 <400> 8
    ggatatcggg tcttcctctt tgcggg 26

```

```

60 <210> 9
    <211> 85
    <212> ADN
    <213> Chimeric Sequence

```

<220>
 <221> gene
 5 <222> (1)..(85)
 <223> Sequence # 9.
 Synthetic fragment coding for 3B protein
 carrying T by C and G by C nucleotide
 substutions, respectively.
 10

<400> 9
 tccagctggtt ggggtttatc atggagtgc taagcccaa caagtgatta aattggatgc 60
 15 agatccagta gagtctcagt tgact 85

<210> 10
 <211> 89
 <212> ADN
 20 <213> Chimeric Sequence

<220>
 25 <221> gene
 <222> (1)..(89)
 <223> Sequence # 10.
 Synthetic fragment coding for 3B protein
 carrying T by C and G by C nucleotide
 substutions, respectively (complementary chain).
 30

<400> 10
 ctagagtcaa ctgagactct actggatctg catccaattt aatcacttgt ttgggcttag 60
 35 tcactccatg ataaaccca acagctgga 89

<210> 11
 <211> 25
 <212> ADN
 40 <213> Chimeric Sequence

<220>
 <221> primer_bind
 <222> (1)..(25)
 45 <223> Sequence # 11.
 Sequence of the oligonucleotide # 11 used
 for the amplification of 3C coding sequence by PCR

50 <400> 11
 tctcagtcaa ctctagaaat agcag 25

<210> 12
 55 <211> 21
 <212> ADN
 <213> Chimeric Sequence

<220>
 60 <221> primer_bind
 <222> (1)..(21)
 <223> Sequence # 12.

Sequence of the oligonucleotide # 12 used for
the amplification of 3C coding sequence by PCR

5 <400> 12
ataagcttga tcaattttct t 21

<210> 13
10 <211> 978
<212> ADN
<213> Hepatitis A virus

<220>
15 <221> gene
<222> Complement((1)..(978))
<223> Sequence # 13.
Sequence corresponding to the region of 3ABC
polyprotein with proteolytic activity having
20 the selfprocessing sites mutated.

<400> 13
25 gaattcctgc agcccgggg atccatggga atttcagatg atgacaatga tagtgcagta 60
gctgagtttt tccggtcttt tccatctggt gaaccatcaa attccaagtt atctagtttt 120
ttccaagctg tcaactaatca caagtgggtt gctgtgggag ctgcagttgg tattcttgga 180
ttgctagtgg gaggatgggt tgtgtataag catTTTTCCC gcaaagagga agaaccaatt 240
ccagctgttg ggtttatca tggagtgact aagcccaaac aagtgattaa attggatgca 300
gatccagtag agtctcagtt gactctagaa atagcaggat tagttaggaa aaatttggtt 360
30 cagtttggag ttggtgagaa aaatggatgt gtgagatggg tcatgaatgc cttaggagtg 420
aaggatgatt ggttgtagt accttctcat gcttataaat ttgaaaagga ttatgaaatg 480
atggagtttt atttcaatag aggtggaact tactattcaa tttcagctgg taatgttggt 540
attcaatctt tagatgtggg attccaagat gttgttctaa tgaaggttcc tacaattccc 600
aagtttagag atattactca acattttatt aagaaaggag atgtgcctag agccttgaat 660
35 cgcttggcaa cattagtgc aaccgttaat ggaactccta tgtaatttc tgagggacct 720
ttaaaaatgg aagaaaaagc cacttatgtt cataagaaga atgatgttac tacggttgat 780
ttgactgtag atcaggcatg gagaggaaaa ggtgaaggtc ttcttggaat gtgtgggtgg 840
gccctagtgt catcaaatca gtccatacaa aatgcaattt tgggtattca tgttgctgga 900
ggaaattcaa ttcttgtggc aaagttgatt actcaagaaa tgtttcaaaa cattgataag 960
40 aaaattgaaa tcaagctt 978

<210> 14
45 <211> 3489
<212> ADN
<213> Hepatitis A virus

<220>
50 <221> gene
<222> Complement((1)..(3489))
<223> Sequence # 14.
Nucleotide sequence CODING for the new
modified open reading frame (ORFm) of the
Cuban M2 strain.

55
60 <400> 14
atgaatatgt ccaaacaagg aattttccag actggttggga gtggccttga ccacatcctg 60
tccttggcag atattgagga agagcaaatg attcagtcctg ttgataggac tgcagtgcact 120
ggagcttctt atttcacttc tgtggaccaa tcttcagttc atactgctga ggttggctca 180
caccaaattg aacctttgaa aacctctgtt gataaacctg gttctaagaa aactcagggg 240
gagaagtttt tcttgattca ttctgctgat tggctcacta cacatgctct ctttcatgaa 300

	gttgcaaaaat	tggatgtggt	gaaactgctg	tacaatgagc	agtttgccgt	ccaaggtttg	360
	ttgagatacc	atacttatgc	aagatttggc	attgagattc	aagttcagat	aaatcccaca	420
	ccctttcagc	aaggaggact	aatctgtgcc	atggttcctg	gtgaccaaag	ttatggttca	480
	atagcatcct	tgactgttta	tcctcatggg	ctgttaaatt	gcaatatcaa	caatgtagtt	540
5	agaataaagg	ttccatttat	ttatactaga	ggtgcttata	attttaaaga	tccacagtac	600
	ccagtttggg	aattgacaat	cagagtttgg	tcagagttga	atattggaac	aggaacctca	660
	gcttatactt	cactcaatgt	tttagctagg	tttacagatt	tggagttgca	tggattaact	720
	cctcttttcta	cacagatgat	gagaaatgaa	tttagagtta	gtactactga	aaatgttgta	780
	aattttgtcaa	attatgaaga	tgcaagggca	aaaatgtctt	ttgctttgga	tcaggaagat	840
10	tggaagtctg	atccttccca	aggtgggtgga	attaaaatta	ctcatttcac	tacctggaca	900
	tccattccaa	ccttagctgc	tcagtttcca	ttcaatgctt	cagattcagt	tgggcaacaa	960
	attaaagtta	taccagtggc	ccatactttt	ttccagatga	caaacactaa	tcctgatcaa	1020
	aaatgtataa	cagccttggc	ctctattttgt	cagatgttct	gcttttgagg	gggagatcct	1080
	gttttcgatt	tccaggtttt	tccaaccaa	tatcattcag	gtaggctgtt	gttttgtttt	1140
15	gttcctggga	atgagttaat	agatgttact	ggaattacat	taaaacaggc	aactactgct	1200
	ccttgtgcag	tgatggacat	tacaggagtg	cagtcaacct	tgagatttcg	tgttccttgg	1260
	atttctgata	caccttatcg	agtgaatagg	tacacgaagt	cagcacatca	aaaaggtgag	1320
	tatactgcc	ttgggaagct	tattgtgtat	tgttataata	gattgacttc	tccttcta	1380
	gttgcttctc	atgtagagtg	taatgtttat	ctttcagcaa	ttaatttggg	atgttttgct	1440
20	cctcttttacc	atgctatgga	tgttaccaca	caggttgagg	atgattcagg	aggtttctca	1500
	acaacagttt	ctcacagaga	gaatgttctc	gatccccaag	ttggcataac	aacctagagg	1560
	gatttaaaag	ggaaagccaa	taggggaaag	atggatgtat	caggagtgc	ggtacctgtg	1620
	ggagctatta	caacaattga	ggatccagtt	ttagcaaaga	aagtacctga	gacatttcct	1680
	gaattgaagc	ctggagaatc	cagacataca	tcagatcaca	tgtctattta	taaattcatg	1740
25	ggaaggtctc	atttcttgtg	tacttttact	tttaattcaa	acaataaaga	gtaacacatt	1800
	ccaataaactc	tgtcttcgac	ttctaatacct	cctcatgggt	taccatcaac	attaaggtgg	1860
	ttctttaatt	tgtttcagtt	gtatagagga	ccattggatt	tgacaattat	aatcacagga	1920
	gccactgatg	tggatgggat	ggcctgggtt	actccagtg	gccttgctgt	cgacacccct	1980
	tggttggaag	agaagtcagc	tttgtctatt	gattataaaa	ctgcccttgg	agctgttaga	2040
30	tttaatacaa	gaagaacagg	gaacattcag	attagattgc	catggtattc	ttatttgtag	2100
	gccgtgtctg	gagcactgga	tggtctggga	gataagacag	attctacatt	tggattgggt	2160
	tctattcaga	ttgcaaatta	caatcattct	gatgaatatt	tgtcctttag	ttgttatttg	2220
	tctgtcacag	agcaatcaga	gttctatttc	cctagagctc	cattaaattc	aaatgctatg	2280
	ttgtccactg	agtccatgat	gagtagaatt	gcagctggag	acttgagtc	atcagtggat	2340
35	gatcccagat	cagaggagga	cagaagattt	gagagtcata	tagaatgtag	gaaaccatat	2400
	aaagaattga	gactggagg	tgggaaacaa	agaatcaaat	atgctcagga	agagttatca	2460
	aatgaagtgc	ttccacctcc	taggaaaatg	aaggggttat	tttcacaagc	tgaattcctg	2520
	cagcccgggg	gatccatggg	aatttcagat	gatgacaatg	atagtgcagt	agctgagttt	2580
	ttccggtctt	ttccatctgg	tgaaccatca	aattccaagt	tatctagttt	tttccaagct	2640
40	gtcactaatc	acaagtgggt	tgctgtggga	gctgcagttg	gtattccttg	attgctagt	2700
	ggaggatggt	tttgtataa	gcatttttcc	cgcaaagagg	aagaaccaat	tccagctgtt	2760
	ggggtttatc	atggagtgc	taagcccaa	caagtgatta	aattggatgc	agatccagta	2820
	gagtctcagt	tgactctaga	aatagcagga	ttagtttagga	aaaatttggt	tcagtttggg	2880
	gttggtgaga	aaaatggatg	tgtgagatgg	gtcatgaatg	ccttaggagt	gaaggatgat	2940
45	tggttgtag	taccttctca	tgcttataaa	tttgaaaagg	attatgaaat	gatggagttt	3000
	tatttcaata	gaggtggaac	ttactattca	atttcagctg	gtaatgttgt	tattcaatct	3060
	ttagatgtgg	gattccaaga	tggtgttcta	atgaagggtc	ctacaattcc	caagtttaga	3120
	gatattactc	aacattttat	taagaaaagg	gatgtgccta	gagccttgaa	tcgcttggca	3180
	acattagtga	caaccgttaa	tggaaactcct	atgttaattt	ctgaggggacc	tttaaaaatg	3240
50	gaagaaaaag	ccacttatgt	tcataagaag	aatgatggta	ctacggttga	tttgactgta	3300
	gatcaggcat	ggagaggaaa	aggtgaaggt	cttcctggaa	tgtgtggtgg	ggccctagt	3360
	tcatcaaatc	agtccatata	aaatgcaatt	ttgggtattc	atgttgctgg	aggaaattca	3420
	attcttgtgg	caaagttgat	tactcaagaa	atgtttcaaa	acattgataa	gaaaattgaa	3480
55	atcaagctt						3489

<210> 15

<211> 51

<212> ADN

60 <213> Chimeric Sequence

<220>

<221> gene
 <222> (1)..(51)
 <223> Sequence # 15.
 Synthetic fragment that reverts the
 transcription start of the vp2 protein.

<400> 15
 gggatggata ttgaggaaga gcaaattgatt cagtccgttg ataggactgc a 51

<210> 16
 <211> 47
 <212> ADN
 <213> Chimeric Sequence

<220>
 <221> gene
 <222> (1)..(47)
 <223> Sequence # 16.
 Synthetic fragment that reverts the transcription
 start of the vp2 protein (complementary chain).

<400> 16
 gtccatcaaa cggactgaat catttgctct tccatcaatat ccatccc 47

<210> 17
 <211> 3426
 <212> ADN
 <213> Hepatitis A virus

<220>
 <221> gene
 <222> Complement((1)..(3426))
 <223> Sequence # 17.
 Sequence coding for the modified open reading
 frame (Δ ORFm) of the Cuban M2 strain of the HAV.
 This sequence does not have the gene coding
 for the VP4 protein.

<400> 17
 gggatggata ttgaggaaga gcaaattgatt cagtccgttg ataggactgc agtgactgga 60
 gcttcttatt tcacttctgt ggaccaatct tcagttcata ctgctgaggt tggctcacac 120
 caaattgaac ctttgaaaac ctctgttgat aaacctggtt ctaagaaaac tcagggggag 180
 aagtttttct tgattcattc tgctgattgg ctactacac atgctctctt tcatgaagtt 240
 gcaaaattgg atgtggtgaa actgctgtac aatgagcagt ttgccgtoca aggtttgttg 300
 agataccata cttatgcaag atttggcatt gagattcaag ttcagataaa tcccacacc 360
 tttcagcaag gaggactaat ctgtgccatg gttcctggtg accaaagtta tggttcaata 420
 gcatccttga ctgtttatcc tcatggctctg ttaaattgca atatcaacaa tgtagttaga 480
 ataaagggttc catttattta tactagaggt gcttatcatt ttaaagatcc acagtaccca 540
 gtttggaagt tgacaatcag agtttggtca gagttgaata ttggaacagg aacctcagct 600
 tatacttcac tcaatgtttt agctagggtt acagatttgg agttgcatgg attaactcct 660
 ctttctacac agatgatgag aaatgaattt agagttagta ctactgaaaa tgttgtaaatt 720
 ttgtcaaatt atgaagatgc aagggcaaaa atgtcttttg ctttgatca ggaagattgg 780
 aagttctgac cttcccaagg tgggtgaatt aaaattactc atttactac ctggacatcc 840
 attccaacct tagctgctca gtttccattc aatgcttcag attcagttgg gcaacaaatt 900
 aaagttatag cagtggaccc atactttttc cagatgacaa acactaatcc tgatcaaaaa 960
 tgtataacag ccttggcctc tatttgtcag atgttctgct tttggagggg agatcttggt 1020
 ttcgatttcc aggtttttcc aaccaaatat cattcaggta ggctgttggt ttgttttggt 1080
 cctgggaatg agttaataga tgttactgga attacattaa aacaggcaac tactgtcctt 1140
 tgtgcagtga tggacattac aggagtgcag tcaaccttga gatttcgtgt tccttggatt 1200

```

tctgatacac cctatcgagt gaataggtac acgaagtcag cacatcaaaa aggtgagtat 1260
actgccattg ggaagcctat tgtgtattgt tataaatagat tgacttctcc ttctaattgtt 1320
gcttctcatg ttagagttaa tgtttatctt tcagcaatta atttggaatg ttttgctcct 1380
ctttaccatg ctatggatgt taccacacag gttggagatg attcaggagg tttctcaaca 1440
5 acagtttcta cagagcagaa tgttctgat ccccaagttg gcataacaac catgagggat 1500
ttaaaggga aagccaatag gggaaagatg gatgtatcag gagtgcaggt acctgtggga 1560
gctattacaa caattgagga tccagtttta gcaaagaaag tacctgagac atttcctgaa 1620
ttgaagcctg gagaatccag acatacatca gatcacatgt ctatttataa attcatggga 1680
aggtctcatt tcttgtgtac ttttactttt aattcaaaca ataaagagta cacatttcca 1740
10 ataactctgt cttcgacttc taatcctcct catggtttac catcaacatt aagggtggtc 1800
tttaatttgt ttcagttgta tagaggacca ttggatttga caattataat cacaggagcc 1860
actgatgtgg atggtatggc ctggtttact ccagtgggcc ttgctgtcga cacccttgg 1920
gtggaaaaga agtcagcttt gtctattgat tataaaactg cccttgagc tgttagattt 1980
aatacaagaa gaacaggga cattcagatt agattgccat ggtattctta tttgtatgcc 2040
15 gtgtctggag cactggatgg cttgggagat aagacagatt ctacatttgg attggtttct 2100
attcagattg caaattacaa tcattctgat gaatatttgt cctttagttg ttatttgtct 2160
gtcacagagc aatcagagtt ctatttccct agagctccat taaattcaaa tgctatgttg 2220
tccactgagt ccatgatgag tagaattgca gctggagact tggagtcac agtggtatgat 2280
cccagatcag aggaggacag aagatttgag agtcatatag aatgtaggaa accatataaa 2340
20 gaattgagac tggaggttgg gaaacaaaga atcaaatatg ctcaggaaga gttatcaaat 2400
aagtgcttc cactcctag gaaaatgaag gggttattt cacaagctga attcctgcag 2460
ccgggggat ccatgggaat ttcagatgat gacaatgata gtgcagtagc tgagtttttc 2520
cggtcttttc catctggtga accatcaa atccaagttat ctagtttttt ccaagctgtc 2580
actaatcaca agtgggttgc tgtgggagct gcagttggta ttcttggatt gctagtggga 2640
25 ggatggtttg tgtataagca tttttccgc aaagaggaag aaccaattcc agctgttggg 2700
gtttatcatg gagtgactaa gcccaaacia gtgattaaat tggatgcaga tccagtagag 2760
tctcagttga ctctagaaat agcaggatta gttaggaaaa atttggttca gtttgagatt 2820
ggtgagaaaa atggatgtgt gagatgggtc atgaatgcct taggagtga ggatgattgg 2880
20 ttgttagtac cttctcatgc ttataaattt gaaaaggatt atgaaatgat ggagttttat 2940
ttcaatagag gtggaactta ctattcaatt tcagctggta atgttgttat tcaatcttta 3000
gatgtgggat tccaagatgt tgttcta atggttccta caattcccaa gtttagagat 3060
attactcaac attttattaa gaaaggagat gtgcctagag ccttgaatcg cttggcaaca 3120
ttagtgacaa ccgttaatgg aactcctatg ttaatttctg agggacctt aaaaatggaa 3180
35 gaaaaagcca cttatgttca taagaagaat gatggtacta cggttgattt gactgtagat 3240
caggcatgga gaggaaaagg tgaaggtctt cctggaatgt gtggtggggc ctagtgtca 3300
tcaaatcagt ccatacaaaa tgcaattttg ggtattcatg ttgctggagg aaattcaatt 3360
cttgtggcaa agttgattac tcaagaaatg tttcaaaaca ttgataagaa aattgaaatc 3420
aagctt 3426

```

40

```

<210> 18
<211> 19
<212> ADN
<213> Chimeric Sequence

```

45

```

<220>
<221> sig_peptide
<222> (1)..(19)
<223> Sequence #18.

```

50

```

Synthetic fragment corresponding to the
KDEL endoplasmic reticulum retention signal
sequence.

```

400> 18

55 atcaaggatg aattgtaat

19

210> 19

211> 21

212> ADN

213> Chimeric Sequence

<220>
 <221> sig_peptide
 <222> (1)..(21)
 <223> Sequence #19.
 5 Synthetic fragment corresponding to the KDEL
 endoplasmic reticulum retention signal sequence.

 <400> 19
 10 cgattacaat tcatccttga t 21

 <210> 20
 <211> 55
 <212> ADN
 15 <213> Chimeric Sequence

 <220>
 <221> D_segment
 <222> (1)..(54)
 20 <223> Sequence # 20
 synthetic fragment modifying the 3' end of
 the 2A protein and introduces a space-bar
 between this one and the KDEL signal.

 25 <400> 20
 cctaggaaaa tgaaggggtt atatgcttct ggaggtgaat tcgatatcaa ggatg 55

 <210> 21
 30 <211> 54
 <212> ADN
 <213> Chimeric Sequence

 <220>
 35 <221> D_segment
 <222> (1)..(54)
 <223> Sequence # 21.
 synthetic fragment modifying the 3' end of
 the 2A protein and introduces a space-bar
 40 between this one and the KDEL signal.

 <400> 21
 aattcatcct tgatatcgaa ttcacctcca gaagcatata accccttcat ttcc 54

 45 <210> 22
 <211> 2555
 <212> ADN
 <213> Hepatitis A virus
 50
 <220>
 <221> gene
 <222> Complement((1)..(2555))
 <223> Sequence # 22.
 55 sequence coding for the structural P1-2A
 proteins joined to endoplasmic reticulum
 retention signal.

 <400> 22
 60 atgaatatgt ccaaacaagg aattttccag actgttggga gtggccttga ccacatcctg 60
 tccttggcag atattgagga agagcaaatg attcagtcg ttgataggac tgcagtgact 120
 ggagcttctt atttcacttc tgtggaccaa tottcagttc atactgctga ggttggtca 180

```

caccaaattg aacctttgaa aacctctgtt gataaacctg gttctaagaa aactcagggg 240
gagaagtttt tcttgattca ttctgctgat tggctcacta cacatgctct ctttcatgaa 300
gttgcaaaat tggatgtggt gaaactgctg tacaatgagc agtttgccgt ccaagggttg 360
ttgagatacc atacttatgc aagatttggc attgagattc aagttcagat aaatcccaca 420
5 ccccttcagc aaggaggact aatctgtgcc atgggttcctg gtgaccaaag ttatggttca 480
atagcatcct tgactgttta tcctcatggg ctgttaaatt gcaatatcaa caatgtagtt 540
agaataaagg ttccatttat ttatactaga ggtgcttatt attttaaaga tccacagtac 600
ccagtttggg aattgacaat cagagtttgg tcagagttga atattggaac aggaacctca 660
gcttataact cactcaatgt tttagctagg tttacagatt tggagttgca tggattaact 720
10 cctcttttcta cacagatgat gagaaatgaa tttagagtta gtactactga aaatgttgta 780
aatttgtcaa attatgaaga tgcaagggca aaaatgtctt ttgctttgga tcaggaagat 840
tgggaagtctg atccttccca aggtggtgga attaaaatta ctcatctcac tacttgga 900
tccattccaa ccttagctgc tcagtttcca ttcaatgctt cagattcagt tgggcaacaa 960
attaaagtta taccagtggg cccatacttt ttccagatga caaacactaa tcctgatcaa 1020
15 aaatgtataa cagccttggc ctctatttct cagatgttct gcttttggag gggagatctt 1080
gttttcgatt tccaggtttt tccaaccaa tatcattcag gtaggctgtt gttttgtttt 1140
gttcctggga atgagttaat agatgttact ggaattacat taaaacaggc aactactgct 1200
ccttgtgcag tgatggacat tacaggagtg cagtcaacct tgagatttcg tgttccttgg 1260
atctctgata caccctatcg agtgaatagg tacacgaagt cagcacatca aaaaggtagg 1320
20 tatactgcca ttgggaagct tattgtgtat tgttataata gattgacttc tccttctaata 1380
cttgcttctc atgttagagt taatgtttat ctttcagcaa ttaatttggg atgttttctc 1440
cctctttacc atgctatgga tgttaccaca caggttggag atgattcagg aggttttctc 1500
acaacagttt ctacagagca gaatgttctt gatccccaag ttggcataac aaccatgagg 1560
gatttaaaag ggaaagccaa taggggaaag atggatgtat caggagtgcg ggtacctgtg 1620
25 ggagctatta caacaattga ggatccagtt ttagcaaaga aagtacctga gacatttcct 1680
gaattgaagc ctggagaatc cagacataca tcagatcaca tgtctattta taaattcatg 1740
ggaaggtctc atttcttgtg tacttttact ttaattcaa acaataaaga gtacacattt 1800
ccaataactc tgtcttcgac ttctaatcct cctcatgggt taccatcaac attaaggtagg 1860
30 ttctttaatt tgtttcagtt gtatagagga ccattggatt tgacaattat aatcacagga 1920
gccactgatg tggatggtat ggcctggttt actccagtgg gccttgctgt cgacaccctt 1980
tggttggaag agaagtcagc tttgtctatt gattataaaa ctgcccttgg agctgttaga 2040
tttaatacaa gaagaacagg gaacattcag attagattgc catggtattc ttatttgtat 2100
gccgtgtctg gagcactgga tggcttggga gataagacag attctacatt tggattggtt 2160
35 tctattcaga ttgcaaatta caatcattct gatgaatatt tgccttttag ttgttatttg 2220
tctgtcacag agcaatcaga gttctatttc cctagagctc cattaaattc aaatgctatg 2280
ttgtccactg agtccatgat gagtagaatt gcagctggag acttggagtc atcagtggat 2340
gatcccgatg cagaggagga cagaagattt gagagtcata tagaatgtag gaaaccatat 2400
aaagaattga gactggaggt tgggaaacaa agaatacaat atgctcagga agagttatca 2460
aatgaagtgc ttccacctcc taggaaatg aaggggttat atgcttctgg aggtgaattc 2520
40 gatatcaagg atgaattgta atcgataccg tcgac 2555

```

<210> 23

<211> 1012

45 <212> ADN

<213> Hepatitis A virus

<220>

<221> gene

50 <222> Complement((1)..(1012))

<223> Sequence # 23.

Sequence coding for the 3ABC polyprotein
and for the endoplasmic reticulum
retention signal.

55

<400> 23

```

gaattcctgc agcccggggg atccatggga atttcagatg atgacaatga tagtgtagta 60
gctgagtttt tccggtcttt tccatctggt gaaccatcaa attccaagtt atctagtttt 120
ttccaagctg tcaactaatca caagtgggtt gctgtgggag ctgcagttgg tattcttggg 180
60 ttgctagtgg gaggatgggt tgtgtataag cttttttccc gcaaagagga agaaccaatt 240
ccagctgttg ggttttatca tggagtgact aagcccaaac aagtgattaa attggatgca 300
gatccagtag agtctcagtt gactctagaa atagcaggat tagttaggaa aaatttgggt 360

```

```

cagtttggag ttggtgagaa aaatggatgt gtgagatggg tcatgaatgc cttaggagt 420
aaggatgatt ggttgtagt accttctcat gcttataaat ttgaaaagga ttatgaaatg 480
atggagtttt atttcaatag aggtggaact tactattcaa ttccagctgg taatgttgtt 540
attcaatctt tagatgtggg attccaagat gttgttctaa tgaaggttcc tacaattccc 600
5 aagtttagag atattactca acattttatt aagaaaggag atgtgcctag agccttgaat 660
cgcttggcaa cattagtgc aaccgttaat ggaactccta tgtaatttc tgagggacct 720
ttaaaaatgg aagaaaaagc cacttatgtt cataagaaga atgatggtac tacggttgat 780
ttgactgtag atcaggcatg gagaggaaaa ggtgaaggtc ttcctggaat gtgtggtggg 840
gccctagtgt catcaaatca gtccatacaa aatgcaattt tgggtattca tggtgctgga 900
10 ggaaattcaa ttcttgtggc aaagttgatt actcaagaaa tgtttcaaaa cattgataag 960
aaaattgaaa tcaagcttcg acctcgaatc aaggatgaat tgtaatcgat ac 1012

<210> 24
15 <211> 2492
    <212> ADN
    <213> Hepatitis A virus

<220>
20 <221> gene
    <222> Complement((1)..(2492))
    <223> Sequence # 24
        Sequence coding for the structural region,
        whitout the vp4 protein, fused to endoplasmic
25        reticulum retention signal.

<400> 24
gggatggata ttgaggaaga gcaaatgatt cagtccgttg ataggactgc agtgactgga 60
gcttcttatt tcacttctgt ggaccaatct tcagttcata ctgctgaggt tggctcacac 120
30 caaattgaac ctttgaaaac ctctgttgat aaacctgggt ctaagaaaac tcagggggag 180
aagtttttct tgattcattc tgctgattgg ctactacac atgctctctt tcatgaagtt 240
gcaaaattgg atgtggtgaa actgctgtac aatgagcagt ttgccgtcca aggtttgttg 300
agataccata cttatgcaag atttggcatt gagattcaag ttcagataaa tcccacaccc 360
tttcagcaag gaggactaat ctgtgccatg gttcctgggtg accaaagtta tggttcaata 420
35 gcatccttga ctgtttatcc tcatggctctg ttaaattgca atatcaacaa tgtagttaga 480
ataaagggttc catttattta tactagaggt gcttatcatt ttaaagatcc acagtaccca 540
gtttgggaat tgacaatcag agtttggcca gagttgaata ttggaacagg aacctcagct 600
tatacttcac tcaatgtttt agctaggttt acagatttgg agttgcatgg attaatcctt 660
ctttctacac agatgatgag aaatgaattt agagttagta ctactgaaaa tgttgtaaat 720
40 ttgtcaaatt atgaagatgc aaggggcaaaa atgtcttttg ctttggatca ggaagatttg 780
aagtctgac cttcccaggg tgggtgaatt aaaattactc atttcactac ctggacatcc 840
attccaacct tagctgtca gtttccattc aatgcttcag attcagttgg gcaacaaatt 900
aaagttatac cagtggacc ctaactttttc cagatgacaa aactaatcc tgatcaaaaa 960
tgtataacag ccttggcctc tatttgtcag atgttctgct tttggagggg agatcttgtt 1020
45 ttcgatttcc aggtttttcc aaccaaatat cattcaggta ggctgttgtt ttgttttgtt 1080
cctgggaatg agttaataga tgttactgga attacattaa aacaggcaac tactgtcctt 1140
tgtgcagtga tggacattac aggagtgcag tcaaccttga gatttcgtgt tccttggatt 1200
tctgatacac cctatcgagt gaataggtac acgaagtcag cacatcaaaa aggtgagtat 1260
actgccattg ggaagcttat tgtgtattgt tataatagat tgacttctcc ttctaattgtt 1320
50 gcttctcatg tttagagttaa tgtttatctt tcagcaatta atttggaaatg ttttgcctct 1380
ctttaccatg ctatggatgt taccacacag gttggagatg attcaggagg tttctcaaca 1440
acagtttcta cagagcagaa tgttcctgat cccaagttg gcataacaac catgagggat 1500
ttaaaaggga aagccaatag gggaaagatg gatgtatcag gagtgcagg acctgtggga 1560
55 gctattacaa caattgagga tccagtttta gcaaagaaag tacctgagac atttcctgaa 1620
ttgaagcctg gagaatccag acatacatca gatcacatgt ctatttataa attcatggga 1680
aggtctcatt tcttgtgtac ttttactttt aattcaaaca ataaagagta cacatttcca 1740
ataaactctg cttcgacttc taatcctcct catggtttac catcaacatt aaggtggttc 1800
tttaatttgt ttcagttgta tagaggacca ttggatttga caattataat cacaggagcc 1860
actgatgtgg atggtatggc ctggtttact ccagtgggcc ttgctgtcga cacccttggg 1920
60 gtggaaaaga agtcagcttt gtctattgat tataaaactg cccttggagc tgtagatttt 1980
aatacaagaa gaacagggaa cattcagatt agattgccat ggtattctta tttgtatgcc 2040
gtgtctggag cactggatgg cttgggagat aagacagatt ctacatttgg attggtttct 2100

```

attcagattg caaattacaa tcattctgat gaatatttgt cctttagttg ttatttgtct 2160
 gtcacagagc aatcagagtt ctatttcctt agagctccat taaattcaaa tgctatgttg 2220
 tccactgagt ccatgatgag tagaattgca gctggagact tggagtcac agtggatgat 2280
 5 cccagatcag aggaggacag aagatttgag agtcatatag aatgtaggaa accatataaa 2340
 gaattgagac tggaggttgg gaaacaaaga atcaaatatg ctcaggaaga gttatcaaat 2400
 gaagtgcctc cacctcctag gaaaatgaag gggttatatg cttctggagg tgaattcgat 2460
 atcaaggatg aattgtaatc gataccgtcg ac 2492